## COLORADO SCHOOL ASTHMA CARE PLAN

COLORADO SCHOOL ASTRIVIA CARE PLAIN	
PARENT/GUARDIAN to complete this portion and sign competed form.	
Name:	Birth date:
Grade:	Parent/Guardian:
Cell Phone:	Home Phone:
Work Phone:	Other Contact:
Phone:	Preferred Hospital:
School:	Teacher:
Triggers: Weather (cold air, wind) Illness E	Exercise Smoke Dog/Cat Dust Pollen
Life threatening allergy: Specify	
Medication Location: school office student possession at all times other location (list)	
If there is no quick relief inhaler at school:	
<ul> <li>Call parents/guardians to pick up student and/or bring inhaler/ medications to school</li> <li>Inform them that if they cannot get to school, 911 may be called</li> </ul>	
I give permission for school personnel to share this information, follow this plan, administer medication and care for my child and, if necessary, contact our physician. I assume full responsibility for providing the school with prescribed medication and delivery/monitoring devices. I approve this Asthma Care Plan for my child.	
PARENT SIGNATURE	
	all items, SIGN and DATE completed form.
GREEN ZONE: Student can do usual activities but should avoid triggers. Asthma is well controlled.	
Pretreatment for strenuous activity: Not Requ	
Pretreatment for strenuous activity: Routinely OR Upon request Explain:(weather, viral, seasonal, other) Give 2 puffs of quick relief med (Check One) Albuterol Nopenex Other: Repeat in 4 hours if needed for additional or ongoing physical activity. If currently experiencing symptoms, follow yellow zone.	
YELLOW ZONE: SICK – UNCONTROLLED ASTH	НМА
IF YOU SEE THIS:	DO THIS:
, 3	1. Stop physical activity
3	2. GIVE QUICK RELIEF MED: (Check One) Albuterol Xopenex Other:  2 puffs Via spacer With mask Other:
<ul><li>Frequent cough</li></ul>	I I / NIITTS I I VIA SNACER I I WITH MASK I I I ITHER.
Complaint of short tightness	
Complains of chest tightness     Unable to tolerate regular activities but	If symptoms do not improve in 10-15 minutes, repeat quick relief medication.
<ul> <li>Unable to tolerate regular activities but</li> </ul>	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> </ul>
<ul> <li>Unable to tolerate regular activities but still talking in complete sentences</li> </ul>	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> <li>Stay with student and maintain sitting position.</li> </ul>
<ul> <li>Unable to tolerate regular activities but still talking in complete sentences</li> <li>Peak flow between and</li> </ul>	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> </ul>
<ul> <li>Unable to tolerate regular activities but still talking in complete sentences</li> <li>Peak flow between and</li> <li>Other:</li> </ul>	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> <li>Stay with student and maintain sitting position.</li> <li>Student may resume normal activities once feeling better.</li> </ul>
<ul> <li>Unable to tolerate regular activities but still talking in complete sentences</li> <li>Peak flow between and</li> <li>Other:</li> </ul>	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> <li>Stay with student and maintain sitting position.</li> <li>Student may resume normal activities once feeling better.</li> <li>If symptoms do not improve in 10-15 minutes or worsen following quick relief med, follow RED ZONE plan.</li> </ul>
■ Unable to tolerate regular activities but still talking in complete sentences ■ Peak flow between and 3 ■ Other: 3  RED ZONE: EMERGENCY SITUATION — SEVER IF YOU SEE THIS: [	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> <li>Stay with student and maintain sitting position.</li> <li>Student may resume normal activities once feeling better.</li> <li>If symptoms do not improve in 10-15 minutes or worsen following quick relief med, follow RED ZONE plan.</li> <li>E ASTHMA SYMPTOMS</li> <li>DO THIS IMMEDIATELY:</li> </ul>
<ul> <li>Unable to tolerate regular activities but still talking in complete sentences</li> <li>Peak flow between and</li> <li>Other:</li> <li>RED ZONE: EMERGENCY SITUATION – SEVERIFYOU SEE THIS:</li> <li>Coughs constantly</li> </ul>	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> <li>Stay with student and maintain sitting position.</li> <li>Student may resume normal activities once feeling better.</li> <li>If symptoms do not improve in 10-15 minutes or worsen following quick relief med, follow RED ZONE plan.</li> <li>E ASTHMA SYMPTOMS</li> <li>DO THIS IMMEDIATELY:</li> <li>GIVE QUICK RELIEF MED: (Check One): Albuterol Xopenex Other:</li> </ul>
<ul> <li>Unable to tolerate regular activities but still talking in complete sentences</li> <li>Peak flow between and</li> <li>Other:</li> <li>RED ZONE: EMERGENCY SITUATION – SEVERIFYOU SEE THIS:</li> <li>Coughs constantly</li> <li>Struggles or gasps for breath</li> </ul>	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> <li>Stay with student and maintain sitting position.</li> <li>Student may resume normal activities once feeling better.</li> <li>If symptoms do not improve in 10-15 minutes or worsen following quick relief med, follow RED ZONE plan.</li> <li>E ASTHMA SYMPTOMS</li> <li>DO THIS IMMEDIATELY:</li> <li>GIVE QUICK RELIEF MED: (Check One): Albuterol Xopenex Other:</li> <li>2 puffs Via spacer With mask Other:</li> </ul>
■ Unable to tolerate regular activities but still talking in complete sentences ■ Peak flow between and # ■ Other: #  RED ZONE: EMERGENCY SITUATION — SEVERIFYOU SEE THIS: # ■ Coughs constantly # ■ Struggles or gasps for breath # ■ Trouble talking (only speaks 3-5 words)	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> <li>Stay with student and maintain sitting position.</li> <li>Student may resume normal activities once feeling better.</li> <li>If symptoms do not improve in 10-15 minutes or worsen following quick relief med, follow RED ZONE plan.</li> <li>E ASTHMA SYMPTOMS</li> <li>DO THIS IMMEDIATELY:</li> <li>GIVE QUICK RELIEF MED: (Check One): Albuterol Xopenex Other:         <ul> <li>2 puffs</li> <li>Via spacer</li> <li>With mask</li> <li>Other:</li> <li>Refer to anaphylaxis plan if student has life threatening allergy.</li> </ul> </li> </ul>
■ Unable to tolerate regular activities but still talking in complete sentences ■ Peak flow between and # ■ Other: *  RED ZONE: EMERGENCY SITUATION — SEVER #  IF YOU SEE THIS:  # ■ Coughs constantly # ■ Struggles or gasps for breath # ■ Trouble talking (only speaks 3-5 words) # ■ Skin of chest and/or neck pull in with # ■ 2	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> <li>Stay with student and maintain sitting position.</li> <li>Student may resume normal activities once feeling better.</li> <li>If symptoms do not improve in 10-15 minutes or worsen following quick relief med, follow RED ZONE plan.</li> <li>E ASTHMA SYMPTOMS</li> <li>DO THIS IMMEDIATELY:</li> <li>GIVE QUICK RELIEF MED: (Check One): Albuterol Xopenex Other:         <ul> <li>2 puffs</li> <li>Via spacer</li> <li>With mask</li> <li>Other:</li> <li>Refer to anaphylaxis plan if student has life threatening allergy.</li> </ul> </li> <li>Call 911 and inform EMS the reason for the call.</li> </ul>
■ Unable to tolerate regular activities but still talking in complete sentences ■ Peak flow between and	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> <li>Stay with student and maintain sitting position.</li> <li>Student may resume normal activities once feeling better.</li> <li>If symptoms do not improve in 10-15 minutes or worsen following quick relief med, follow RED ZONE plan.</li> <li>E ASTHMA SYMPTOMS</li> <li>THIS IMMEDIATELY:</li> <li>GIVE QUICK RELIEF MED: (Check One): Albuterol Xopenex Other:         <ul> <li>2 puffs</li> <li>Via spacer</li> <li>With mask</li> <li>Other:</li> <li>Refer to anaphylaxis plan if student has life threatening allergy.</li> </ul> </li> <li>Call 911 and inform EMS the reason for the call.</li> <li>Call parents/guardians and school nurse.</li> </ul>
■ Unable to tolerate regular activities but still talking in complete sentences ■ Peak flow between and # ■ Other: #  RED ZONE: EMERGENCY SITUATION — SEVER #  IF YOU SEE THIS:	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> <li>Stay with student and maintain sitting position.</li> <li>Student may resume normal activities once feeling better.</li> <li>If symptoms do not improve in 10-15 minutes or worsen following quick relief med, follow RED ZONE plan.</li> <li>E ASTHMA SYMPTOMS</li> <li>THIS IMMEDIATELY:</li> <li>GIVE QUICK RELIEF MED: (Check One): Albuterol Xopenex Other:         <ul> <li>2 puffs</li> <li>Via spacer</li> <li>With mask</li> <li>Other:</li> <li>Refer to anaphylaxis plan if student has life threatening allergy.</li> </ul> </li> <li>Call 911 and inform EMS the reason for the call.</li> <li>Call parents/guardians and school nurse.</li> <li>Encourage student to take slower deeper breaths.</li> </ul>
■ Unable to tolerate regular activities but still talking in complete sentences ■ Peak flow between and 4 ■ Other: 8  RED ZONE: EMERGENCY SITUATION – SEVERIFYOU SEE THIS:	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> <li>Stay with student and maintain sitting position.</li> <li>Student may resume normal activities once feeling better.</li> <li>If symptoms do not improve in 10-15 minutes or worsen following quick relief med, follow RED ZONE plan.</li> <li>E ASTHMA SYMPTOMS</li> <li>DO THIS IMMEDIATELY:</li> <li>GIVE QUICK RELIEF MED: (Check One): Albuterol Xopenex Other: Quiffs Via spacer With mask Other: Refer to anaphylaxis plan if student has life threatening allergy.</li> <li>Call 911 and inform EMS the reason for the call.</li> <li>Call parents/guardians and school nurse.</li> <li>Encourage student to take slower deeper breaths.</li> <li>Stay with student and remain calm.</li> </ul>
■ Unable to tolerate regular activities but still talking in complete sentences ■ Peak flow between and # ■ Other: #  RED ZONE: EMERGENCY SITUATION – SEVER   IF YOU SEE THIS:	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> <li>Stay with student and maintain sitting position.</li> <li>Student may resume normal activities once feeling better.</li> <li>If symptoms do not improve in 10-15 minutes or worsen following quick relief med, follow RED ZONE plan.</li> <li>E ASTHMA SYMPTOMS</li> <li>THIS IMMEDIATELY:</li> <li>GIVE QUICK RELIEF MED: (Check One): Albuterol Xopenex Other:         <ul> <li>2 puffs</li> <li>Via spacer</li> <li>With mask</li> <li>Other:</li> <li>Refer to anaphylaxis plan if student has life threatening allergy.</li> </ul> </li> <li>Call 911 and inform EMS the reason for the call.</li> <li>Call parents/guardians and school nurse.</li> <li>Encourage student to take slower deeper breaths.</li> </ul>
■ Unable to tolerate regular activities but still talking in complete sentences ■ Peak flow between and	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> <li>Stay with student and maintain sitting position.</li> <li>Student may resume normal activities once feeling better.</li> <li>If symptoms do not improve in 10-15 minutes or worsen following quick relief med, follow RED ZONE plan.</li> <li>E ASTHMA SYMPTOMS</li> <li>DO THIS IMMEDIATELY:</li> <li>GIVE QUICK RELIEF MED: (Check One): Albuterol Xopenex Other: Specific With mask Other: Refer to anaphylaxis plan if student has life threatening allergy.</li> <li>Call 911 and inform EMS the reason for the call.</li> <li>Call parents/guardians and school nurse.</li> <li>Encourage student to take slower deeper breaths.</li> <li>Stay with student and remain calm.</li> <li>If symptoms do not improve, continue to give quick relief medication until EMS arrives.</li> <li>School personnel should not drive student to hospital.</li> </ul>
■ Unable to tolerate regular activities but still talking in complete sentences ■ Peak flow between and	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> <li>Stay with student and maintain sitting position.</li> <li>Student may resume normal activities once feeling better.</li> <li>If symptoms do not improve in 10-15 minutes or worsen following quick relief med, follow RED ZONE plan.</li> <li>E ASTHMA SYMPTOMS</li> <li>DO THIS IMMEDIATELY:</li> <li>GIVE QUICK RELIEF MED: (Check One): Albuterol Xopenex Other: Refer to anaphylaxis plan if student has life threatening allergy.</li> <li>Call 911 and inform EMS the reason for the call.</li> <li>Call parents/guardians and school nurse.</li> <li>Encourage student to take slower deeper breaths.</li> <li>Stay with student and remain calm.</li> <li>If symptoms do not improve, continue to give quick relief medication until EMS arrives.</li> <li>School personnel should not drive student to hospital.</li> </ul> APPROPRIATE BOX(ES) Is medications, and in my opinion, can carry and use his/her inhaler at school independently with Officials after using inhaler.
■ Unable to tolerate regular activities but still talking in complete sentences ■ Peak flow between and 4 ■ Other: 5  RED ZONE: EMERGENCY SITUATION – SEVERI IF YOU SEE THIS:	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> <li>Stay with student and maintain sitting position.</li> <li>Student may resume normal activities once feeling better.</li> <li>If symptoms do not improve in 10-15 minutes or worsen following quick relief med, follow RED ZONE plan.</li> <li>E ASTHMA SYMPTOMS</li> <li>DO THIS IMMEDIATELY:</li> <li>GIVE QUICK RELIEF MED: (Check One): Albuterol Xopenex Other: Refer to anaphylaxis plan if student has life threatening allergy.</li> <li>Call 911 and inform EMS the reason for the call.</li> <li>Call parents/guardians and school nurse.</li> <li>Encourage student to take slower deeper breaths.</li> <li>Stay with student and remain calm.</li> <li>If symptoms do not improve, continue to give quick relief medication until EMS arrives.</li> <li>School personnel should not drive student to hospital.</li> </ul> APPROPRIATE BOX(ES) Is medications, and in my opinion, can carry and use his/her inhaler at school independently with Officials after using inhaler.
■ Unable to tolerate regular activities but still talking in complete sentences ■ Peak flow between and 4 ■ Other: 5  RED ZONE: EMERGENCY SITUATION – SEVERIFYOU SEE THIS:	<ul> <li>If symptoms do not improve in 10-15 minutes, repeat quick relief medication.</li> <li>Call parents/guardians and school nurse.</li> <li>Stay with student and maintain sitting position.</li> <li>Student may resume normal activities once feeling better.</li> <li>If symptoms do not improve in 10-15 minutes or worsen following quick relief med, follow RED ZONE plan.</li> <li>E ASTHMA SYMPTOMS</li> <li>DO THIS IMMEDIATELY:</li> <li>GIVE QUICK RELIEF MED: (Check One): Albuterol Xopenex Other: Refer to anaphylaxis plan if student has life threatening allergy.</li> <li>Call 911 and inform EMS the reason for the call.</li> <li>Call parents/guardians and school nurse.</li> <li>Encourage student to take slower deeper breaths.</li> <li>Stay with student and remain calm.</li> <li>If symptoms do not improve, continue to give quick relief medication until EMS arrives.</li> <li>School personnel should not drive student to hospital.</li> </ul> APPROPRIATE BOX(ES) Is medications, and in my opinion, can carry and use his/her inhaler at school independently with Officials after using inhaler.

Photo of child